

# Process-Oriented Quantitative Evaluation of IPCC CMIP5 Simulations Using A-Train Satellite Observations

Hui Su, Jonathan Jiang, Chengxing Zhai, Vince Perun, Janice T. Shen

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

with Contributions from

Anthony Del Genio, Larissa Nazarenko (GISS/USA); William Rossow (CCNY/USA); Leo J. Donner, L. Horowitz, C. Seman (GFDL/USA), Andrew Gettelman (NCAR/USA); Tongwen Wu (BCC/China); Mark Ringer, Roger Saunders (UKMO/UK); J-L Dufresne (IPSL/France); Jason Cole (CCCMA/Canada); Tristan L'Ecuyer (WISC/USA); Steve Platnick (GSFC/USA); Graeme Stephens, Joao Teixeira, Baijun Tian (JPL/USA)

# Objective

- To use A-Train satellite observations to evaluate CMIP5 simulations of clouds and water vapor and thus contribute to IPCC AR5
  - Cloud feedback remains the largest uncertainty for climate projections.
  - Accurate simulations of current climate is a necessary condition for credible future projections.




Image credit: NASA




Image credit: Wikipedia/NOAA

# What New Capabilities Enabled by A-Train?

1. Vertical profiles of water vapor and clouds
2. Simultaneous multiple measurements enable physics-driven process understanding
3. Knowledge of observational uncertainties enables quantitative assessment


	Spatial Mean	Variance	Correlation
IWC, LWC	$G_n^{IWC, LWC} = \max \left[ 0, 1 - \frac{1}{n_g} \frac{ \ln(m_{mod}^{IWC, LWC}) - \ln(m_{obs}^{IWC, LWC}) }{\ln \epsilon_{n, obs}^{IWC, LWC}} \right]$	$G_v^{IWC, LWC} = \max \left[ 0, 1 - \frac{1}{n_g} \frac{ \ln \sigma_{mod}^{IWC, LWC} - \ln \sigma_{obs}^{IWC, LWC} }{\ln \epsilon_{v, obs}^{IWC, LWC}} \right]$	$G_c = \max [0, C_{mod, obs}]$
H <sub>2</sub> O	$G_n^{H_2O} = \max \left[ 0, 1 - \frac{1}{n_g} \frac{ m_{mod}^{H_2O} - m_{obs}^{H_2O} }{\epsilon_{n, obs}^{H_2O} m_{obs}^{H_2O}} \right]$	$G_v^{H_2O} = \max \left[ 0, 1 - \frac{1}{n_g} \frac{ \sigma_{mod}^{H_2O} - \sigma_{obs}^{H_2O} }{\epsilon_{v, obs}^{H_2O}} \right]$	

- For IWC and LWC, a logarithm of the ratio of modeled value to the observed is used due to large spread among the models.
- $\sigma$  in the denominator represents the uncertainty range of the observation.
- The scaling factor  $n_g = 3$  (=4 for LWC at 900 hPa)
  - A zero G score means: 1) for H<sub>2</sub>O, the model-observation difference is > 3 times the observational uncertainty; 2) for IWC and LWC, the model value is either > 2<sup>3</sup>=8 times (16 times for 900hPa) or < 1/8 (1/16 for 900hPa) of the corresponding observation.

References: Douglass et al. [1999], Waugh and Eyring [2008], and Gettelman et al. [2010]

See Jiang et al. (2011, to be submitted)


# Approach



# Step 1

Performance of current  
climate simulations for  
water vapor and clouds



# Evaluation of Multi-year Climatology



Jiang et al. (2011,  
to be submitted)




# Comparing Spatial Mean, Variance and Correlation



Jiang et al. (2011,  
to be submitted)

# Model Performance





# Overall Score and Rank


AR5 Model	Overall Score	Rank
BCC csm1	0.56	8
BCCR noresm	0.70	3
CCCMA canesm2	0.61	7
CCCMA cm4	0.61	7
GFDL am3	0.64	6
GFDL cm3	0.64	6
GISS e2-h	0.52	9
GISS e2-r	0.51	10
INM cm4	0.49	11
IPSL cm5a	0.66	4
NCAR cam5	0.65	5
UKMO hadgem2-a	0.73	1
UKMO hadgem2-es	0.71	2
Multi-model-mean	0.74	
Weighted multi-model-mean	0.75	

Jiang al. (2011, to be submitted)

## Step 2

Performance of current  
climate simulations for  
water vapor and clouds


processes that govern the  
simulations




```
graph LR; A([processes that govern the simulations]) --> B[Performance of current climate simulations for water vapor and clouds]
```

The diagram illustrates a process flow. It begins with an oval containing the text 'processes that govern the simulations'. A white arrow points from this oval to a rounded rectangle containing the text 'Performance of current climate simulations for water vapor and clouds'. The background is a dark blue gradient with wavy lines at the top.

# Conditional Sampling



# Conditional Sampling



## Step 3


Performance of current  
climate simulations for  
water vapor and clouds




Impact on current climate  
radiation and precipitation




# Model Spread in Cloud Forcing Sensitivity




# Model Spread in Cloud Forcing Sensitivity



# Model Spread in Precipitation Sensitivity



# Correlation with Precipitation



600 hPa

# Summary

- Significant improvements are found from CMIP3 to CMIP5 in simulated IWP and LWP.
- Water vapor is generally better simulated than clouds.
- Model spreads in the upper troposphere are much larger than those in the lower and middle troposphere.
- The simulated relationships of clouds with large-scale dynamic and thermodynamic regimes are drastically different -> large errors in model physics
- Boundary layer clouds constitute the largest spread for the net cloud forcing sensitivity
- Mid-tropospheric clouds constitute the largest spread for the precipitation sensitivity






More work is needed .....


# Back-up Slides

# Relevance to TOA Cloud Forcing




200 hPa

# Model Spread in Cloud Forcing Sensitivity




# Model Spread in Cloud Forcing Sensitivity






# Correlation with Precipitation




100 hPa

# Correlation with Precipitation



215 hPa

# Correlation with Precipitation



900 hPa